

SHEEP Dewormer Chart

Important --Please read notes below before using this chart

1 ml = 1cc	Valbazen (albendazole) <u>ORALLY</u>	SafeGuard (fenbendazole) <u>ORALLY</u>	Ivomec Sheep Drench (ivermectin) <u>ORALLY</u>	Prohibit (levamisole) <u>ORALLY</u>	Cydetin Sheep Drench (moxidectin) <u>ORALLY</u>
Weight Pounds (lbs)	7.5 mg/kg 0.75 ml/ 25 lb	5 mg/kg 0.6 ml/ 25 lb	0.2 mg/kg 2.9 ml/ 25 lb	8 mg/kg 2 ml/ 25 lb	0.2 mg/kg 2.3 ml/25 lb
20	0.6	0.5	2.3	1.5	1.8
25	0.75	0.6	2.9	1.8	2.3
30	0.9	0.7	3.4	2.2	2.7
35	1.1	0.8	4.0	2.6	3.2
40	1.2	0.9	4.5	2.9	3.6
45	1.4	1.0	5.1	3.3	4.1
50	1.5	1.1	5.7	3.7	4.5
55	1.7	1.3	6.2	4.0	5.0
60	1.8	1.4	6.8	4.4	5.4
65	2.0	1.5	7.4	4.7	5.9
70	2.1	1.6	8.0	5.1	6.3
75	2.3	1.7	8.5	5.5	6.8
80	2.4	1.8	9.1	5.8	7.2
85	2.6	1.9	9.7	6.2	7.7
90	2.7	2.0	10.2	6.6	8.1
95	2.9	2.1	10.8	6.9	8.6
100	3.0	2.2	11.4	7.3	9.1
105	3.2	2.3	1.02	7.7	9.5
110	3.3	2.5	12.5	8.0	10
115	3.5	2.6	13.1	8.4	10.5
120	3.6	2.7	13.7	8.8	10.9
125	3.8	2.8	14.2	9.1	11.4
130	3.9	2.9	14.8	9.5	11.8
140	4.2	3.0	15.4	10.2	12.7
150	4.5	3.1	16.0	11.0	13.6

Valbazen Suspension (11.36 % or 113.6 mg/ml): 7.5 mg/kg orally; approved in sheep with meat withdrawal time of 7 days. Do NOT use in pregnant ewes in the first trimester of pregnancy.

Safe-Guard/ Panacur Suspension (10% or 100 mg/ml): Note that SafeGuard is not approved for use in sheep. Sheep dose is 5 mg/kg orally; meat withdrawal time of 6 days.

Ivomec Drench for Sheep (0.08% or 0.8 mg/ml): 0.2 mg/kg orally; approved in sheep with meat withdrawal time of 11 days. Protect from light.

Prohibit Soluble Drench Powder (Sheep): (Note that this drug is also sold as Levasol and Tramsiol) 8 mg/kg ORAL dose. Approved for use in sheep with meat withdrawal of 3 days. Solution prepared by dissolving a 52 gram packet in 1 quart (943 ml) of water. This yields a solution with 49.6 mg/ml. Always make sure to follow directions on packet when preparing.

If dosing lambs, it is safer to dilute further (1 packet in 2 quarts of water), and then administer twice the amount listed on the chart. The larger volume administered will provide a wider margin for safety if there are small errors in dosing.

Cydetin Sheep drench (1 mg/ml): 0.2 mg/kg orally; approved in sheep with meat withdrawal time of 14 days.

NOTE for Guideline for Dewormer (Anthelmintic) Dosages in Sheep

This chart was developed by Ray M. Kaplan, DVM, PhD and Lisa Williamson, DVM, MS (University of Georgia). It is provided as a possible guideline for anthelmintic (deworming) dosages for sheep. Producers should always consult their veterinarian for advice on their specific management situation for determining which dewormer(s) are best to use in their flock, and the proper dosages for their flock. Meat withdrawal times should always be strictly adhered to.

Note that drug resistance in parasites of sheep is extremely common. The effectiveness of a particular dewormer should always be tested before being used by performing either a Fecal Egg Count Reduction Test (FECRT) or DrenchRite larval development assay (contact Dr. Kaplan's laboratory [706-542-0742] for more information about the DrenchRite test).

Updated September 2014